

# Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☒ Coloured covers/  
Couverture de couleur

☐ Coloured pages/  
Pages de couleur

☐ Covers damaged/  
Couverture endommagée

☐ Pages damaged/  
Pages endommagées

☐ Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée

☐ Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées

☐ Cover title missing/  
Le titre de couverture manque

☒ Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées

☐ Coloured maps/  
Cartes géographiques en couleur

☐ Pages detached/  
Pages détachées

☐ Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)

☒ Showthrough/  
Transparence

☐ Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur

☐ Quality of print varies/  
Qualité inégale de l'impression

☒ Bound with other material/  
Relié avec d'autres documents

☐ Continuous pagination/  
Pagination continue

☒ Tight binding may cause shadows or distortion along interior margin/  
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure

☐ Includes index(es)/  
Comprend un (des) index

☐ Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/  
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

Title on header taken from: /  
Le titre de l'en-tête provient:

☐ Title page of issue/  
Page de titre de la livraison

☐ Caption of issue/  
Titre de départ de la livraison

☐ Masthead/  
Générique (périodiques) de la livraison

☒ Additional comments: / Pagination is as follows: p. 63-70.  
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

UNIVERSITY OF TORONTO  
STUDIES

PHYSIOLOGICAL SERIES

No. 12: THE ACTION OF YEAST FRACTIONS ON THE  
GROWTH OF RATS, BY CASIMIR FUNK and ARCHIBALD BRUCE  
MACALLUM

(REPRINTED FROM THE JOURNAL OF BIOLOGICAL CHEMISTRY, VOL. XXVII)

THE UNIVERSITY LIBRARY: PUBLISHED BY  
THE LIBRARIAN, 1916

**University of Toronto Studies**  
**COMMITTEE OF MANAGEMENT**

---

**Chairman: ROBERT ALEXANDER FALCONER, M.A., LL.D., D.D.**  
**President of the University**

**PROFESSOR W. J. ALEXANDER, Ph.D.**

**PROFESSOR W. H. ELLIS, M.A., M.B.**

**PROFESSOR J. J. MACKENZIE, B.A., M.B.**

**PROFESSOR J. P. McMURRICH, Ph.D.**

**PROFESSOR G. H. NEEDLER, Ph.D.**

**PROFESSOR GEORGE M. WRONG, M.A.**

**General Editor: H. H. LANGTON, M.A.**

**Librarian of the University**

Recru

By

(From

R

The  
growth  
the  
of be  
for th  
fracti  
experi  
sized  
stabil  
on gro  
dimini  
beri-b  
metho

The  
prom  
the be  
with p  
idue o  
and al  
precipi  
silver  
to their

\* Sen

## STUDIES ON GROWTH.

### IV. THE ACTION OF YEAST FRACTIONS ON THE GROWTH OF RATS.

By CASIMIR FUNK AND ARCHIBALD BRUCE MACALLUM.\*

(From the General Memorial Hospital, Harriman Research Laboratory, Roosevelt Hospital, New York, and the Department of Pathological Chemistry, University of Toronto.)

(Received for publication, August 3, 1916.)

The close relationship existing between the beri-beri and growth problems suggests the possibility of a fractionation of the active substance along lines already used in the investigation of beri-beri. Accordingly phosphotungstic acid was selected for the first attempt to separate out a physiologically active fraction which would stimulate the growth of young rats. The experimental difficulties which have repeatedly been emphasized in the investigation of the beri-beri vitamine, due to instability of this substance, were also encountered in the study on growth. The physiological activity of the different fractions diminishes with each manipulation and both the problems of beri-beri and of growth will not be solved until more suitable methods for the isolation of vitamine are available.

The results obtained so far clearly indicate that the growth-promoting substance is analogous to and possibly identical with the beri-beri vitamine and can be almost entirely precipitated with phosphotungstic acid. Subsequent fractionation of the residue obtained from the decomposed precipitate with silver nitrate and also with silver nitrate and baryta has shown that the precipitate containing purine bases and the filtrate from the silver nitrate and baryta precipitation are entirely negative as to their growth-promoting action; whereas the substances pre-

\* Senior Fellow in Medical Research.

precipitated with silver nitrate and baryta possess traces of the activity of the initial phosphotungstic acid precipitate. The experimental evidence indicates that considerably larger quantities of vitamins are necessary for stimulating growth than for curing beri-beri, and the losses occurring during fractionation are more apparent in the former than in the latter case. However, it must be admitted that while it is uncertain whether these two substances are chemically different, the results obtained do not exclude such a possibility. Lloyd's reagent, as recommended by Seidell,<sup>1</sup> has also been used as a precipitant without much success, as the rats on the filtrate have also shown increments in growth.

In the first instance autolyzed yeast was slightly acidified with hydrochloric acid and completely precipitated with phosphotungstic acid, carefully avoiding an excess of this reagent. After allowing the mixture to stand for 24 hours, the precipitate was collected on a Buchner funnel and repeatedly washed with a cold solution of phosphotungstic acid containing hydrochloric acid. The precipitate was then decomposed by the method described by Van Slyke,<sup>2</sup> with a mixture of amyl alcohol and ether and hydrochloric acid, only a small quantity of the precipitate remaining unchanged. After filtration of this small fraction the aqueous extract was evaporated *in vacuo* and the residue made up to a known volume and mixed in the diet in quantities calculated from the amount of autolyzed yeast necessary to promote growth. However, the quantity of this fraction had to be doubled and even tripled in order to obtain satisfactory results. The phosphotungstic acid filtrate was worked out in a similar way. This process offers the advantage that the yeast fraction is completely freed from substances soluble in lipid solvents. The purine fraction was obtained from the phosphotungstic acid precipitate fraction by precipitation with silver nitrate and subsequent decomposition with sulfuretted hydrogen. The filtrate from the purine bases was precipitated with baryta in the usual way and the precipitate decomposed, freed from traces of baryta, evaporated, and the residue mixed with

<sup>1</sup> Seidell, A., *U. S. Public Health Report*, No. 325, 1916.

<sup>2</sup> Van Slyke, D. D., *J. Biol. Chem.*, 1915, xxii, 281.

the di  
was re  
tate o  
alcohol  
with t  
paper  
nitrate  
further  
stituen  
ize the  
to the  
preven

A lar  
on the  
of only  
Rats 9  
out the  
days to  
fraction  
view th  
precipit

Casein...  
Sugar...  
Starch...  
Lard...  
Agar...  
Salts...  
NaHCO<sub>3</sub>...  
Phosphotungstic acid...



the diet. The filtrate from the fraction containing vitamins was reprecipitated with phosphotungstic acid and the precipitate obtained after thorough washing, decomposed with amyl alcohol and ether. The results with the purine fraction and also with the silver nitrate-baryta filtrate are not included in this paper. They were entirely negative. The effect of the silver nitrate-baryta fraction was not sufficiently marked to encourage further investigation. The diet contained lard as the fat constituent, and 1 per cent sodium bicarbonate was added to neutralize the hydrochloric acid present in this fraction. Orange juice to the extent of 1 cc. a day was added to the drinking water to prevent the onset of scorbutic symptoms.

A large number of rats were kept on the above diets, especially on the phosphotungstic precipitate and filtrate and the records of only a few were selected for publication, in order to save space. Rats 9 and 10 were kept on phosphotungstic acid filtrate throughout the experiment. Rats 11 and 12 were changed after 34 days to the diet containing the phosphotungstic precipitate fraction which was followed by an improvement warranting the view that the growth-promoting substance is contained in this precipitate (Fig. 2 b).

Diets (Gm.).

	1.	2.
Casein.....	22	22
Sugar.....	10	10
Starch.....	23	23
Lard.....	30	30
Agar.....	2	2
Salts.....	2	2
NaHCO <sub>3</sub> .....	1	1
Phosphotungstate precipitate.....		10
"    filtrate.....	10	

Rat 9 (male) and Rat 10 (female).

Rats 11 and 12. Males.

0-96 days Diet 1.

0-34 days Diet 1.

28-100 " 1 cc. orange juice.

35-100 " " 2.

76-100 triple vitamin addition.

Days	Weight		Average daily food intake.	Days	Weight		Average daily food intake.
	9.	10.			11.	12.	
	gm.	gm.	gm.		gm.	gm.	gm.
0	35.2	41.8		0	37.9	34.2	
4	42.8	47.3	9.5	4	46.2	41.8	7.6
8	43.0	47.3	7.8	8	45.8	44.2	5.5
12	44.6	49.3	7.0	12	42.8	45.3	7.6
16	46.8	55.4	8.7	16	45.3	45.8	7.8
20	46.2	56.6	7.5	20	46.9	45.7	7.7
24	46.6	54.7	7.8	24	46.2	46.5	6.6
28	46.5	52.8	5.5	28	44.3	45.0	5.4
32	42.9	49.9	7.4	32	40.4	39.4	4.7
36	37.8	41.6	7.8	36	39.8	37.2	5.6
40	36.8	41.6	7.7	40	50.5	48.2	9.1
44		41.8	6.6	44	52.2	50.0	8.9
48		43.9	5.3	48	54.8	57.0	8.0
52		44.0	5.3	52	58.0	60.7	8.7
56		43.8	3.8	56	63.5	67.3	11.0
60		41.6	3.5	60	66.3	71.0	12.4
64		44.6	3.5	64	66.1	70.3	8.8
68		41.0	3.2	68	70.6	75.7	11.1
72		43.5	3.5	72	75.0	82.6	9.9
76		42.9	3.4	76	76.0	84.8	11.9
80		45.0	3.0	80	77.6	91.1	11.1
84		42.2	3.1	84	75.0	85.2	9.6
88		44.1	2.7	88	79.4	90.0	8.8
92		43.0	3.2	92	78.5	90.9	4.4
96		41.0	2.9	96	74.7	91.7	6.4
100				100	70.0	76.3	6.4

A second series of experiments was carried out on pressed yeast which had been heated with 10 per cent sulfuric acid at 90-95° for 6 hours. The hydrolysate was filtered, diluted with an equal volume of water, and precipitated with phosphotungstic acid. After standing for 24 hours the precipitate was filtered at the pump and well washed with 5 per cent sulfuric acid.

The precipitate was decomposed in the ordinary way with baryta. The final solution, slightly acid, was neutralized with sodium carbonate, carefully avoiding an excess, distilled *in vacuo*, standardized, and definite quantities were added to the diet. The

filtrate of the phosphotungstic acid precipitation was treated in a similar way.

The diet containing the substances precipitated by phosphotungstic acid was fed to four rats (Rats 80 to 83, Fig. 1) and enabled them to double their original weight after 32 to 36 days. This is about double the time required when yeast is the source of vitamins and the depreciation is due to the fractionation with the precipitating reagent.

Two rats (84 and 85, Fig. 2 a) were fed the diet containing the residue from the phosphotungstic filtrate. After 28 days they had added only a third to their original weight and had all the external symptoms of an acute deficiency. Then the diet with the precipitate was substituted and in 11 days they rapidly doubled their original weight and presented a normal appearance.

Diets (Gm.).

	1.	2.
Casein.....	22	22
Sugar.....	10	10
Starch.....	24	24
Fat (lard).....	30	30
Salt.....	6	6
Agar.....	2	2
Residue phosphotungstic acid precipitate equal to.....	6	Dried 6/ yeast.
Residue phosphotungstic acid filtrate equal to.....		

Rats 80, 81, and 82. Males.

Rat 83. Female.

0-36 days Diet 1.

Days.	Weight.		Average daily food intake.	Weight.		Average daily food intake.
	80.	81.		82.	83.	
	gm.	gm.	calories	gm.	gm.	calories
0	24	25		20	30	
4	31	33	29.8	23.5	37	27.8
8	36	40.5	33.5	28	43	32.8
12	36	43	35.7	29	45	33.3
16	38	44.5	40.5	29	48	40.5
20	40	48	28.6	31	51	38.3
24	42	49	30.1	34	54	30.2
28	44.5	52	27.8	38	57	35.0
32	46	54	28.7	40	59	27.4
36	49	56	25.0	42.5	61	29.8






FIG. 1. Rats 80, 81, 82, and 83 were kept on a diet containing decomposed phosphotungstic acid precipitate from yeast.

FIG. 2a. Rats 84 and 85 were kept first on a diet containing the residue from phosphotungstic filtrate of yeast. The marked deficiency was corrected when this addition was changed on the point marked on the chart to the corresponding precipitate.




FIG. 2b. Rats 9 and 10 were kept on a diet containing phosphotungstic acid filtrate throughout the experiment. Rats 11 and 12 were changed to phosphotungstic precipitate at the point marked on the chart, with a marked improvement in growth and general appearance.

Fig 2b.

Rat 84. Female.

Rat 85. Male.

0-28 days Diet 2.

29-39 " " 1.

Days.	Weight.		Average daily food intake.
	84.	85.	
	<i>gm.</i>	<i>gm.</i>	<i>calories.</i>
0	45	45	
4	48	49	28.6
8	53	54.5	46.3
12	55	55	42.9
16	55	55	35.9
20	57	57	50.9
24	58	58	48.6
28	59	61	47.9
32	69	68	54.6
36	84	82	70.3
39	89	90	66.3

## SUMMARY.

The fractionation of yeast with phosphotungstic acid shows that the growth-promoting substance is carried down with the precipitate and a large part of its activity is lost during the fractionation. The instability of this substance when fractionated with silver salts presents greater difficulty than that experienced during the fractionation of the beri-beri vitamin. It seems possible that both these problems can only be solved when more adequate methods are available.



# UNIVERSITY OF TORONTO STUDIES

## PHYSIOLOGICAL SERIES

- |   |      |
|---|------|
| No. 1: The structure, micro-chemistry and development of nerve-cells, with special reference to their nuclein compounds, by F. H. SCOTT ..... | 0.50 |
| No. 2: On the cytology of non-nucleated organisms, by A. B. MACALLUM .....  | 0.75 |
| No. 3: Observations on blood pressure, by R. D. RUDOLF...   | 0.75 |
| No. 4: The chemistry of wheat gluten, by G. G. NASMITH...   | 0.50 |
| No. 5: The palaeochemistry of the ocean, by A. B. MACALLUM  | 0.25 |
| No. 6: The absorption of fat in the intestine, by G. E. WILSON  | 0.50 |
| No. 7: The distribution of fat, chlorides, phosphates, potassium and iron in striated muscle, by MAUD L. MENTEN.....                          | 0.25 |
| No. 8: Surface tension and vital phenomena, by A. B. MACALLUM   | 1.00 |
| No. 9: On the distribution of potassium in renal cells, by C. P. BROWN.....   | 0.25 |
| No. 10: On the probable nature of the substance promoting growth in young animals, by CASIMIR FUNK and A. BRUCE MACALLUM.....                 | 0.25 |
| No. 11: A new conception of the glomerular function, by T. G. BRODIE.....   |      |
| On changes in the glomeruli and tubules of the kidney accompanying activity, by T. G. BRODIE and J. J. MACKENZIE .....                        | 1.00 |